

Version 2\_dated 03 Mar 2022 DSRB Reference: 2021/01024 PECC-GST-ED-2022-1420-v1



#### What You Should Know

We are glad to have you here to start your Hepatitis C treatment journey.

- Yes! Chronic Hepatitis C is a treatable disease with a recovery rate of more than 95%.
- This will require your commitment to take your medication(s) as prescribed by your doctor.



### Hepatitis C

#### What You Should Know



#### **How is Hepatitis C Treated Nowadays?**

- We use all-ORAL medications to achieve recovery.
- In Singapore, there are 2-3 types of Hepatitis C medication that your doctor may choose to treat you with initially.
- There are slight differences in the duration and total pills you need to take daily, based on the treatment choice.
- Either way, as long you can complete the treatment, you have a more than 95% chance to recover from Hepatitis C.

# More than **95% chance** to be healthy again!





#### What You Should Know

**PRESCRIPTION** 

#### **Dos and Don'ts During Treatment**

- Understand the correct way to take your medication.
- Take your medication at the same time every day.
- Do not stop medication yourself or omit any tablets.
- Avoid over-the-counter medication (e.g. antacids) or supplements while on treatment.
- Talk to your doctor before you stop the Hepatitis C treatment or if starting any new medication.

## Hepatitis C

#### What You Should Know







 If you have liver hardening (cirrhosis), your doctor may add another medication called Ribavirin to increase your treatment success rate.

500 813  Ribavirin can cause anaemia, a condition which causes low red blood cells in your body. Your doctor will monitor this.



#### What You Should Know



## **Contraception Is Important When You Are On Hepatitis C Treatment**

- If you do not practise safe sex during the course of treatment, your baby will be at risk of birth defects.
- Use TWO effective contraceptive methods to prevent pregnancy, until six months after you complete Hepatitis C treatment.

### Hepatitis C

#### What You Should Know



## What are some Common Side Effects and Precautions?

Observe for potential drug allergy (for example, rash, itchiness, eye-lid swelling, facial swelling, etc.). Stop the medication when you observe these symptoms, and seek medical attention immediately.

 Otherwise, the side effects are mostly manageable. The common side effects are nausea, headache, fatigue, diarrhoea and insomnia. Common
Side Effects
Nausea
Nausea
Headache
Fatigue
Diarrhoea
Insomnia

Version 2\_dated 03 Mar 202 DSRB Reference: 2021/0102



## Coping with Side Effects

### Hepatitis C

#### Coping with Side Effects



#### Nausea



- Eat smaller but frequent meals
- Avoid oily, spicy, acidic or sweet foods

#### Headache



- Maintain adequate fluid intake and take plenty of rest
- Take paracetamol when it is necessary

#### **Fatigue**



- Maintain adequate fluid intake and take plenty of rest
- Have well-balanced meals
- Exercise regularly

#### Diarrhoea



- Take plenty of fluids e.g. water, clear soup
- Avoid coffee, tea, dairy products or spicy/ acidic foods

#### Insomnia



- Avoid excessive daytime naps
- Avoid coffee or tea before bedtime
- Have a light snack before bedtime
- Avoid eating heavy dinner

## **Anaemia** (if on Ribavirin)



 Talk to your doctor if you feel tired, breathless, giddy or have chest pain


## IT IS NOT NECESSARY TO STOP TAKING YOUR MEDICATIONS

because of side effects,

as these mostly last for a short time and are manageable.

If side effects persist or worsen, inform your doctor.

### Epclusa® (Sofosbuvir/Velpatasvir) +/- Ribavirin



- Take your medications at the same time each day.
- Do NOT stop taking your medications without talking to your doctor.
- If you miss a dose, take the missed dose the same day as soon as you remember; take your next dose at your regular time the next day.
- If you miss more than 1-2 days of doses, call your doctor/clinic.

Version 2\_dated 03 Mar 2022 DSRB Reference: 2021/01024 PECC-GST-ED-2022-1420-v1

#### Maviret® (Glecaprevir/Pibrentasvir)

Start date: Tre	atment duration:
End date: Dat	e to test for cure:
	Morning (C)
☐ Maviret®: Take 3 tablets once daily	Time:

- Take your medications at the same time each day.
- Do NOT stop taking your medications without talking to your doctor.
- If you miss a dose, take the missed dose the same day as soon as you remember; take your next dose at your regular time the next day.
- If you miss more than 1-2 days of doses, call your doctor/clinic.



#### What You Should Know

What if the Hepatitis C treatment was not successful this round?

- Do not give up, as your doctor may discuss with you other treatment options using a different way.
- You still stand a pretty good chance of recovery if you can complete this treatment.



#### Have a question?

The following contact numbers may be helpful to you.





Clinic 4B hotline

Tel: 6889 4445

**Operating hours:** 

Monday to Friday: 8:00 am to 5:30 pm

Saturday, Sunday and

Public Holiday: Closed

• TTSH General Enquiries Contact

Tel: 6256 6011

Email: contact@ttsh.com.sg

• TTSH Level 4 Pharmacy

Tel: 6889 4353

**Operating hours:** 

Monday to Friday: 9:00 am to 6:00 pm

Saturday, Sunday and

Public Holiday: Closed

• TTSH Level B2 Pharmacy contact and operating hours

Tel: 6357 2040

**Operating hours:** 

Monday to Friday:

8:30 am to 6:00 pm

Saturday:

8:30 am to 6:00 pm

Sunday and

**Public Holiday: Closed** 



#### Have a question?

The following contact numbers may be helpful to you.



• Tel: 6440 2444

Email: highpoint\_hwh@hcsa.org.sg



• Emergency ambulance

Tel: 995

• Non-emergency ambulance

Tel: 1777

### Hepatitis C

#### What You Should Know



Last but not least...

- As you already know, Hepatitis C virus is primarily spread through infected blood. For example, when you have raw wounds or during piercings, tattoos, or when sharing used syringes or needles.
- You may get a new infection if you have risk factors again.

 Your health MATTERS for your FUTURE.





#### What You Should Know





Share with your friends and peers what you already know:

- Hepatitis C is a virus transmitted through blood and can harm the liver.
- RISK factors include piercings, tattoos.
- There is a CURE for Hepatitis C today.
- The treatment is affordable.
- The side effects are manageable.

And you can receive help from this programme:





