

# Tips for Managing Diarrhea While on Chemotherapy

## Things you can do on your own



### Keep track of your diarrhea:

- Write down how many bowel movements you have each day

### Stay hydrated and change your diet to lessen your symptoms:

- Drink at least 6-8 cups of clear liquids, such as water, daily while having diarrhea to prevent dehydration
- Avoid caffeine, alcohol, milk
- Avoid greasy, fatty, and fried foods
- Eat small, frequent meals throughout the day instead of 3 big meals
- Drink sports drinks or oral rehydration salt to replenish your electrolytes

### Keep the skin around your anus healthy:

- If your skin is getting irritated after bowel movements, gently wipe yourself with baby wipes and then apply barrier creams or diaper ointments

### Medication:

- Take loperamide 2mg after each loose stools, up to max of 16mg per day

## When to seek medical attention



- If you have a fever  $\geq 38^{\circ}\text{C}$
- If your diarrhea worsens despite following the above tips
- If you are having signs of dehydration. Signs of dehydration include:
  - Having dark yellow or very little urine
  - Having a dry mouth/being thirsty
  - If you have  $\geq 7$  episodes of diarrhea a day
- If there is large amount of watery stoma output (for patients with stoma)
- If there is blood in your stool
- If your stools are black in color, as this can sometimes be a sign of internal bleeding
- If there is severe abdominal pain or persistent abdominal pain



Scan the QR Code with your smart phone to access the information online or visit <https://nhghealth.com.sg>

Department of Oncology  
PECC-ONC-ED-2025-2082-v1

© Tan Tock Seng Hospital, Singapore 2025. All rights reserved. All information correct as of September 2025. No part of this document may be reproduced, copied, reverse compiled, adapted, distributed, commercially exploited, displayed or stored in a database, retrieval system or transmitted in any form without prior permission of Tan Tock Seng Hospital. All information and material found in this document are for purposes of information only and are not meant to substitute any advice provided by your own physician or other medical professionals.