

Tips for Managing Mouth or Throat Sores While on Chemotherapy

Things you can do on your own



- Use a toothbrush with extra soft bristles, and toothpastes for sensitive gums
- Rinse mouth regularly with alcohol free mouth wash or salt water
- Avoid harsh mouthwashes that contain alcohol
- If you are using dentures, rinse and brush the dentures after eating and soak them in denture solution overnight
- Use lip balm to keep your lips moist
- Eat foods that are cold, moist, and soft
- Avoid foods that are salty, spicy, hard, or crunchy
- Drink liquids through a straw
- You may try using topical medications such as choline salicylate (e.g. bonjela) or triamcinolone (e.g. oracort E) or painkillers (e.g. paracetamol) to relieve the pain

REMINDER - Always keep your care team updated about over-the-counter medicines you are taking. If your care team tells you they are not safe for you, do not take them.

When to seek medical attention



- If you have signs of an infection or fever $\geq 38^{\circ}\text{C}$
- If your mouth pain is severe
- If you are unable to eat or drink or have minimal intake for more than 12 hours
- If you feel dizzy or lightheaded when you stand or have reduced urine output



Scan the QR Code with your smart phone to access the information online or visit <https://nhghealth.com.sg>

Department of Oncology
PECC-ONC-ED-2025-2091-v1

© Tan Tock Seng Hospital, Singapore 2025. All rights reserved. All information correct as of September 2025. No part of this document may be reproduced, copied, reverse compiled, adapted, distributed, commercially exploited, displayed or stored in a database, retrieval system or transmitted in any form without prior permission of Tan Tock Seng Hospital. All information and material found in this document are for purposes of information only and are not meant to substitute any advice provided by your own physician or other medical professionals.