

PHARMACY

Drug Information

Medications to Treat Heart Failure



What is Heart Failure?

- Heart failure (HF) happens when the heart cannot pump enough blood to the rest of the body. This results in insufficient oxygen being delivered for normal body functions, especially when one is active or exercising.
- The most common cause of HF is coronary heart disease, which is the narrowing of blood vessels that supply blood and oxygen to heart muscle.
- Other causes include high blood pressure, abnormal heart rhythms, thyroid disorders, severe anemia and infection.

What are the Medications for Heart Failure and the Side Effects?

Medications and how They Work	Common Side Effects	Management and Advice
<p>Angiotensin-Converting Enzyme Inhibitors (ACE-I)</p> <ul style="list-style-type: none"> • <i>Captopril</i> • <i>Enalapril</i> • <i>Lisinopril</i> • <i>Perindopril</i> • <i>Ramipril</i> <p>Angiotensin II Receptor Blockers (ARB)</p> <ul style="list-style-type: none"> • <i>Candesartan</i> • <i>Losartan</i> • <i>Valsartan</i> <p>Angiotensin Receptor-Nepriylsin Inhibitor (ARNI)</p> <ul style="list-style-type: none"> • <i>Valsartan-Sacubitril</i> <p><u>How do they work</u> Prevent further weakening of the heart resulting from heart failure. Also reduce workload of the heart and blood pressure.</p>	<ul style="list-style-type: none"> • Dizziness • Dry cough • (usually ACE-I) • May cause high Potassium level: dizziness, muscle weakness or fast heart beat • Rare side effects: <ul style="list-style-type: none"> ○ Swelling of the hands, face, lips, eyes, tongue, or throat ○ Breathing difficulty ○ Swallowing difficulty 	<ul style="list-style-type: none"> • Get up slowly from sitting or lying position to minimize dizziness • Inform your doctor if you have persistent dry cough, severe dizziness or symptoms of high Potassium levels occur • Inform your doctor if you intend to get pregnant or are currently pregnant. • Seek medical attention immediately if you experience any rare side effects <p>Note: ACE-I and ARNI should not be taken within 36 hours of each other should your doctor switch you from one medication to another.</p>

Medications and how They Work	Common Side Effects	Management and Advice
Beta Blockers <ul style="list-style-type: none"> • <i>Bisoprolol</i> • <i>Carvedilol</i> • <i>Metoprolol</i> • <i>Nebivolol</i> <u>How do they work</u> Reduce heart rate and blood pressure, thereby reducing workload of the heart and improve heart function over time.	<ul style="list-style-type: none"> • Dizziness • Tiredness • Cold hands and feet • Breathlessness • Sexual dysfunction 	<ul style="list-style-type: none"> • Get up slowly from a sitting or lying position to minimize dizziness • Symptoms such as tiredness, cold hands and feet may resolve over time • Inform your doctor if breathlessness or persistent giddiness occur • Do not stop taking this medicine without discussing with your doctor as it may worsen your condition
Mineralocorticoid Receptor Antagonist <ul style="list-style-type: none"> • <i>Spironolactone</i> • <i>Eplerenone</i> <u>How do they work</u> Suppress the hormone (aldosterone) that affects the heart, thereby maintaining or improving heart function.	<ul style="list-style-type: none"> • Breast tenderness or enlargement (Specific to Spironolactone, especially in males) • Menstrual changes • May cause high Potassium level: dizziness, muscle weakness or fast heart beat 	<ul style="list-style-type: none"> • Take after meals for better absorption • Get up slowly from sitting or lying position to minimize dizziness • Inform your doctor if severe dizziness, breast swelling/ pain, menstrual changes, or symptoms of high Potassium levels occur
I _f Channel Inhibitor <ul style="list-style-type: none"> • <i>Ivabradine</i> <u>How does it work</u> Lowers the heart rate thereby reduces the oxygen demand and workload of the heart.	<ul style="list-style-type: none"> • Dizziness • Sensation of enhanced brightness • Blurred vision 	<ul style="list-style-type: none"> • Inform your doctor if persistent dizziness or changes in vision occur

Medications and how They Work	Common Side Effects	Management and Advice
Diuretics (“Water pills”) <i>Thiazide Diuretics</i> <ul style="list-style-type: none"> • <i>Metolazone</i> <i>Loop Diuretics</i> <ul style="list-style-type: none"> • <i>Bumetanide</i> • <i>Frusamide</i> <u>How do they work</u> Reduce symptoms of breathlessness, abdominal bloatedness and leg swelling by increasing urine output.	<ul style="list-style-type: none"> • Nausea, vomiting • Dizziness • May cause low Potassium levels: dizziness, muscle weakness, muscle cramp or fast heart beat 	<ul style="list-style-type: none"> • Take in the morning or late afternoon to avoid having to wake up at night to pass urine • Frequent urination is expected. Inform your doctor if you are unable to pass urine • Inform your doctor if you experience symptoms of low Potassium levels • Your doctor may prescribe you with Potassium Chloride supplements. Inform your doctor if you cannot swallow the Potassium tablets whole
Arterial Vasodilators <ul style="list-style-type: none"> • <i>Hydralazine</i> <u>How do they work</u> Relax blood vessels to reduce blood pressure (afterload).	<ul style="list-style-type: none"> • Dizziness, headache • Rare side effects: <ul style="list-style-type: none"> ○ Fast heartbeat ○ Leg swelling ○ Drug-induced lupus: fever, rashes, muscle or joint pain ○ Blood disorders: Mouth ulcers, fatigue, fever, sore throat or unexplained bleeding or bruises 	<ul style="list-style-type: none"> • Rise slowly from sitting or lying position to minimize dizziness • Seek medical attention immediately if any of the rare side effects occurs

Medications and how They Work	Common Side Effects	Management and Advice
Venous Vasodilators (Nitrates) <ul style="list-style-type: none"> • <i>Isosorbide Dinitrate</i> • <i>Isosorbide-5-Mononitrate CR (controlled release)</i> <u>How do they work</u> Relax blood vessels and reduces the workload of the heart (preload).	<ul style="list-style-type: none"> • Dizziness • Diarrhea • Flushing of face 	<ul style="list-style-type: none"> • Isosorbide-5-Mononitrate CR should be swallowed whole, do not crush or chew the tablets • Headache and flushing of face usually improve with time • Inform your doctor if severe headache or persistent dizziness occur • Do not take with Phosphodiesterase-5 Enzyme Inhibitors [e.g. Sildenafil (Viagra®), Tadalafil (Cialis®), Vardenafil (Levitra®)] without discussing with your doctor
Cardiac Glycosides <ul style="list-style-type: none"> • <i>Digoxin</i> <u>How does it work</u> Improve the pumping action of the heart and improve exercise tolerance. They can also be used to control irregular fast heart rate.	<ul style="list-style-type: none"> • Dizziness • Diarrhea, nausea, vomiting • Abdominal pain • Symptoms of digoxin toxicity (rare) include: <ul style="list-style-type: none"> ○ Persistent diarrhea, nausea, vomiting ○ Visual changes (such as yellow or blurred vision, seeing bright spots) 	<ul style="list-style-type: none"> • Take after meals to reduce nausea and vomiting • Seek medical attention immediately if any of the Digoxin toxicity symptoms occur

Medications and how They Work	Common Side Effects	Management and Advice
Calcium Channel Blockers <i>Dihydropyridine derivatives:</i> <ul style="list-style-type: none"> • <i>Amlodipine</i> • <i>Nifedipine LA (Long Acting)</i> <u>How do they work</u> Relax blood vessels and reduces blood pressure.	<ul style="list-style-type: none"> • Dizziness • Headache • Flushing of face • Lower leg swelling 	<ul style="list-style-type: none"> • Nifedipine LA (Long Acting) should be swallowed whole, do not crush or chew the tablets. • Rise slowly from sitting or lying position to minimize dizziness • Inform your doctor if dizziness is persistent • Elevating your legs may help to reduce leg swelling
Anti-platelets (" Blood Thinner ") <ul style="list-style-type: none"> • <i>Aspirin</i> • <i>Ticagrelor</i> • <i>Clopidogrel</i> • <i>Prasugrel</i> <u>How do they work</u> Prevent the formation or growth of blood clots and reduce the chance of heart attack or stroke.	<ul style="list-style-type: none"> • Gastric irritation (For Aspirin) • Easy bruising or bleeding 	<ul style="list-style-type: none"> • Take after meals to reduce gastric irritation • Be careful to avoid injury • Avoid rubbing bruises and apply pressure to small cuts to stop bleeding • See a doctor immediately if unusual bleeding/bruising, or black and tarry stools occurs • Inform your doctor or dentist that you are on a blood thinner before any surgical or dental procedures

Medications and how They Work	Common Side Effects	Management and Advice
<p>Cholesterol-Lowering Agents (“Statins”)</p> <ul style="list-style-type: none"> • Atorvastatin • Simvastatin • Rosuvastatin • Lovastatin • Pravastatin <p><u>How do they work</u> Decrease bad cholesterol (LDL) and the fatty acids in the blood (triglyceride), hence reducing risks of heart attack.</p>	<ul style="list-style-type: none"> • Muscle ache • Rare side effects <ul style="list-style-type: none"> ○ Unexplained severe or generalized muscle pain/weakness ○ Tea-coloured (Dark brown) urine ○ Yellowing of skin or whites of eyes ○ Severe abdominal pain 	<ul style="list-style-type: none"> • Take Lovastatin after food for better absorption • Take Simvastatin or Lovastatin at night • Seek medical attention immediately if you experience any of the rare side effects

If you develop rashes, facial swelling or shortness of breath after taking the medication, you could be allergic to the medication. Please seek medical attention immediately.

How are the Medications Taken?

- If you miss a dose, take the missed dose as soon as you remember. If it is almost time for your next dose, take only the usual dose. Do not double the dosage.

Can the Medications be Taken with Other Medications?

- Please also inform your doctor or pharmacist if you are taking other medications, including supplements, traditional or herbal remedies.

Do I Need Regular Monitoring When Taking These Medicines?

- Regular monitoring of blood tests may be required.

Are There any Special Precautions That I Need to Take?

- It is important to inform your doctor if you intend to conceive or breastfeed while taking these medications.
- If you experience dizziness while taking more than one of the medications, you can try staggering the dosing time of the medications at intervals of 2 hours.

What Else can I do to Help My Condition?

- Limit your water and salt intake as recommended, to prevent excessive retention of water.
- Weigh yourself at the same time every day after emptying your bladder in the morning. Inform your doctor if sudden weight gain of more than 1kg in one day or more than 3kg in a week occurs.
- Exercise at least 3 times a week for 30 minutes each time.
- Stop smoking and avoid alcohol.

How do I Store These Medications?

- Store your medication in a cool, dry place away from heat, moisture and direct sunlight.

If you have any problem with your treatment, please contact your doctor, pharmacist or nurse clinician.

Please keep all medications out of reach of children.

