

Agomelatine

Medication Information Leaflet

What is this medication for?

Agomelatine belongs to a class of medications known as Melatonin Agonist. It is used to treat symptoms of depression (low mood) such as:

- Feeling sad
- Feeling hopeless or useless
- Feeling bad about yourself
- Thoughts of ending your life
- Poor sleep, energy or appetite

Agomelatine may be used for other conditions. Check with your healthcare professional if you are unsure why you are given this medication.

How should I take/use this medication?

- Follow the instructions on your medication label and take this medication as prescribed by your doctor.
- This medication may be taken with or without food. It is usually taken before sleeping.
- This medication needs to be taken regularly for a few weeks before you feel its full benefits. You should not stop taking this medication even if you do not feel better at the beginning. Continue to take this medication even after you feel better to prevent your symptoms from returning.
- Do check with your doctor on how long you will have to be on this medication.
- Do not stop taking your medication without checking with your healthcare professional.

What should I do if I forget to take/use this medication?

If you forget to take a dose, take it as soon as you remember. However, if it is more than half of the time to the next dose, skip the missed dose and take the next dose at your usual timing. Do not take two doses to make up for the missed dose.

What precautions should I take?

Inform your healthcare professional if:

- You are allergic to this medication or any of the other ingredients of this medication
- You are pregnant, planning to become pregnant, or breastfeeding. If you are pregnant and/or breastfeeding, your doctor would have discussed the potential benefits and

side-effects with you and should you have further questions or concerns, do consult your doctor.

- You have or have ever had:
 - Other drug allergies
 - Kidney or liver disease
 - Other mental health conditions such as bipolar disorder or mania (abnormally high or excited mood)

Your mental condition may change unexpectedly, especially in the first few weeks of treatment and during dose changes. Check with your doctor immediately if you experience the following:

- New or increasing thoughts of harming yourself
- Worsening anxiety, restlessness, violent behaviour, or
- Other changes in mood or behaviour.

Be sure that your family or caregiver are also aware of the above symptoms so they can call your doctor if you are unable to seek treatment on your own.

What are some side-effects that I need to seek medical advice immediately?

The symptoms of a drug allergy include one or more of the following:

- Swollen face/eyes/lips/tongue
- Difficulty in breathing
- Itchy skin rashes over your whole body

Very rarely (affect less than 1% of patients) this medication may cause:

- Feelings of intense agitation or anxiety, restlessness or inability to sit still
- Talking, feeling, and acting with excitement that you cannot control
- Yellowing of skin or eyes, pain in your upper stomach or dark-coloured urine
- Severe weakness or dizziness
- Blurred or double vision
- Unusual behaviour or thoughts of hurting yourself or others

If you experience any of these symptoms, you should stop your medication and consult your healthcare professional immediately.

What are some other side-effects of this medication and what can you do about it?

The mentioned side effects generally affect up to 10% of patients.

- Drowsiness or feeling tired
 - Avoid driving, or taking part in activities that requires concentration.
- Nausea or vomiting
 - Take the medication with or after food. Take small but frequent meals and avoid fatty or spicy food

- Diarrhoea
 - Drink plenty of water to replace the loss of fluid
- Constipation (difficulty passing motion)
 - Drink more water (if you do not have any fluid restriction), eat more high- fibre foods, and exercise regularly
 - Consult your doctor or pharmacist if you need medication to relieve your condition.
- Dizziness
 - Get up slowly from sitting or lying down position
- Mild nervousness
 - May occur at the start of treatment and improves over time
- Headache, increased sweating, back pain or difficulty in sleeping
 - Speak to your doctor if any of these affect your daily life

Most of the side effects listed here will improve with time. Speak with your doctor if they continue, get worse or are affecting your daily life.

What food or medication should I avoid when I take this medication?

Avoid drinking alcohol while taking this medication as it may make you feel very drowsy. This medication may interact with other medications. If you intend to take any over the counter medication, supplements, herbal remedy or traditional Chinese medicine, consult your doctor or healthcare professional first to confirm that it is safe to take with Agomelatine.

How should I store this medication?

Store in a cool and dry place, away from direct sunlight. Keep this medication away from children.

How should I throw away this medication safely?

Pack this medication into a trash bag and seal it tightly before throwing into the rubbish chute or bin.

Disclaimers

If you take more than the recommended dose, please seek medical advice immediately. The information provided on this page does not replace information from your healthcare professional. Please consult your healthcare professional for more information.

This article is jointly developed by members of the National Medication Information workgroup. The workgroup consists of cluster partners (National Healthcare Group, National University Health System and SingHealth), community

pharmacies (Guardian, Unity and Watsons) and Pharmaceutical Society of Singapore.

Last updated on June 2019