

PHARMACY

Drug Information

Atezolizumab (Tecentriq®) Injection



What is Atezolizumab?

- Atezolizumab is used to treat cancer by enhancing your immune system to attack the cancer cells.

How is Atezolizumab Administered?

- Atezolizumab is given as an injection into the vein (intravenous) every three weeks. The first dose is given over 60 minutes and subsequent infusions can be given over 30 minutes if previous doses are well tolerated.

Can Atezolizumab be Administered With Other Medications?

- Please inform your doctor or pharmacist if you are taking other medications, including Traditional Chinese Medicines, Jamu medicines or supplements.

What are the Important Side Effects of Atezolizumab?

Infusion related reaction

- These are usually mild or moderate, but rarely, they can be more severe. You may be given drugs before the infusion to reduce the chance of a reaction happening.
- If you do have an infusion related reaction, this can usually be treated by slowing or stopping the drip until the reaction is over. Any future infusions will be given at a slower rate.
- *Inform the nurses immediately during infusion of Atezolizumab if you experience the following:*
 - Flu-like symptoms, such as headache, feeling flushed, having a fever, chills or dizziness
 - Red, warm and itchy bumps on the skin
 - Swelling of the lips, tongue or throat
 - Breathlessness, wheezing, a cough or sudden difficulty breathing
 - Chest tightness or chest pain

- Medications prescribed by your doctor should be beneficial for your condition even though they may result in some inevitable side effects.

Side Effect	Management
Fatigue or Tiredness	<ul style="list-style-type: none"> • Do not drive or operate machinery if you are feeling tired. Inform your doctor if fatigue is persistent or worsens.
Diarrhoea (colitis, inflammation of the bowel)	<ul style="list-style-type: none"> • Your doctor may prescribe medicines to control diarrhoea • Please proceed to the A&E department if you experience abdominal pain, diarrhoea, cramping, mucus or blood in stool, or dark and sticky stools occur
Nausea or vomiting	<ul style="list-style-type: none"> • Take small, frequent meals. • Avoid smells that precipitate nausea. • Take medicines for nausea or vomiting prescribed by your doctor. • Avoid tight-fitting clothes around the waist.
Changes in the way the liver works (Hepatitis, inflammation of the liver)	<ul style="list-style-type: none"> • Please proceed to the A & E department immediately if you experience any yellowing of eye white and skin or tea-coloured urine, pale stools or severe abdominal pain.
Lung problems (Pneumonitis, inflammation of the lung)	<ul style="list-style-type: none"> • Please proceed to the A&E department immediately if you experience difficulty in breathing, shortness of breath, chest pain and/or new or worsening cough

Side Effect	Management
Hormone abnormalities (inflammation of the pituitary, thyroid, pancreas and adrenal glands can occur and affect the production of certain hormones)	<ul style="list-style-type: none"> • Some hormone levels can be monitored with periodic blood tests • Some of the signs and symptoms include headaches, nausea, vomiting, constipation, rapid heart rate, increased sweating, extreme fatigue, weakness, changes in your voice, changes in memory and concentration, increased hunger or thirst, increased urination, weight gain, hair loss, dizziness, feeling cold all the time, and changes in mood or behavior • Inform your doctor should any of these occur
Brain and/or nerve problems	<ul style="list-style-type: none"> • Inform your doctor if you experience headache, drooping of eyelids, double vision, trouble swallowing, weakness of arms, legs or face, or numbness or tingling in the hands or feet.
Changes in eyesight	<ul style="list-style-type: none"> • Please inform your doctor if you experience eye pain, swelling, redness, blurred vision, and/or sensitivity to light or other changes.
Pancreatitis	<ul style="list-style-type: none"> • Inform your doctor if you are experiencing bloating, indigestion, fatty stools, loss of appetite, sweating, abdominal pain, and weight loss

If you develop rashes, facial swelling or shortness of breath after taking the medication, you could be allergic to the medication. Please seek medical attention immediately.

Are There any Special Precautions That I Need to Take?

- It is important to inform your doctor if you intend to conceive or breastfeed while taking this medication.
- You must not become pregnant during treatment with Atezolizumab and must use an effective method of contraception. You should take appropriate contraceptive measures during and up to 5 months after stopping treatment. If pregnancy occurs during your treatment, you must inform your doctor immediately.
- Atezolizumab may have an anti-fertility effect, which could be irreversible. You should discuss with your doctor regarding fertility preservation prior to starting treatment.
- You should not breast-feed during treatment.

What Else can I do to Help My Condition?

- Exercise
Light exercise is recommended as it helps to improve mobility, balance and even emotional well-being. Avoid strenuous exercise, and rest when you are feeling tired.
- Stay positive
Understand your condition and take an active role in managing it. Do not be afraid to talk about it with your doctor/ pharmacist/ family members. Repressed feelings can be harmful to your immediate well-being and long-term health.
- Adequate hydration
Maintain adequate hydration with 2 to 3 liters (8 to 12 cups) of fluids per day unless informed by your doctor to restrict fluids. This will decrease your chances of getting constipation and also prevent dehydration.
- Medical and dental treatment
If you need to go into hospital for any reason other than cancer, always inform the doctors and nurses that you are having chemotherapy. Talk to your cancer doctor if you think you need dental treatment. Always inform your dentist that you are having chemotherapy

If you have any problem with your treatment, please contact your doctor, pharmacist or nurse clinician.

You can also find useful information with regards to your medication on this website:

<https://www.macmillan.org.uk/information-and-support>



© Tan Tock Seng Hospital, Singapore 2021. All rights reserved. All information correct as of August 2021. No part of this document may be reproduced, copied, reverse compiled, adapted, distributed, commercially exploited, displayed or stored in a database, retrieval system or transmitted in any form without prior permission of Tan Tock Seng Hospital. All information and material found in this document are for purposes of information only and are not meant to substitute any advice provided by your own physician or other medical professionals.

PECC-PHA-ED-2021-1273-v1