

Brinzolamide Eye Drop

Medication Information Leaflet

What is this medication for?

Brinzolamide belongs to a class of medications called carbonic anhydrase inhibitors, which is used to treat glaucoma. Glaucoma is a condition where the nerve in the eye is damaged, and is usually associated with increased eye pressure. Untreated glaucoma can affect your eyesight and cause blindness. Brinzolamide helps to lower the pressure in your eye by reducing production of fluid in the eye.

How should I take/use this medication?

- Proper use of eye drop
 - Wash your hands with soap and water, and read the instructions on the label.
 - Shake the bottle.
 - Tilt your head backwards and look up.
 - Gently pull the lower eyelid down to form a pocket and squeeze the bottle to apply one drop into your eye. To prevent the tip of the bottle from getting dirty, make sure the tip does not touch the eye or any other surface.
 - Close your eyes gently and wipe off any excess eye drops around your eyes with a clean tissue.
 - Keep your eyes closed and press down gently on the inner corner of your eye for 1 to 2 minutes to allow the medication to be absorbed.
 - If you need to use more than 1 drop of medication, or if you are using another type of eye drop, wait at least 5 to 10 minutes before applying the next eye drop. This is to prevent the first drop of medication from being washed away by the second drop before it has time to work. This also prevents any medication interactions between different eye drops.
 - If your doctor prescribed both eye drops and eye ointments, you should use the eye drops first.
 - Click on this link for a step-by-step guide on how to apply eye drops
<https://www.healthhub.sg/a-z/medications/364/A-Step-By-Step-Guide-On-How-To-Instil-Eye-Drops>
- For contact lens wearers
 - Remove your contact lens before each application of medication.
 - When using hard lens, you can put them on directly after applying the eye drops.

- When using soft lens, you can put them on 30 minutes after applying the eye drops. Otherwise, your lens may be damaged by the preservatives present in the eye drops.
- If you need to apply eye drops twice a day, it is recommended that you apply them once in the morning before you put on your lens, and at the end of the day after you remove your lens.
- Do not stop using your medication without checking with your healthcare professional.

What should I do if I forget to take/use this medication?

If you forget to use a dose, use it as soon as you remember. Then use your next dose at the usual time. Do not use two doses to make up for the missed dose.

What precautions should I take?

Inform your healthcare professional if:

- You are allergic to this medication or any of the other ingredients of this medication.
- You are pregnant, planning to become pregnant, or breastfeeding .
- You are taking any other medications, including supplements, traditional medications and herbal remedies.

What are some common side-effects of this medication?

- Stinging or burning sensation
- Redness
- Itchiness
- Eye irritation
- Taste disturbance (bitter or unusual taste in the mouth)
 - After instilling the eye drop, close your eyes and press down gently on the inner corner of the eye for 1 to 2 minutes. It may help to reduce this side effect.
- Blurred vision
 - If you experience blurred vision, do not drive or operate any machinery until your vision is clear.

If these side effects happen and do not go away, check with your healthcare professional as soon as possible.

What are some rare but serious side-effects that I need to seek medical advice immediately?

The symptoms of a drug allergy include one or more of the following:

- Swollen face/eyes/lips/tongue
- Difficulty in breathing
- Itchy skin rashes around your eyes or over your whole body

If you experience any of these symptoms, you should stop your medication and see your healthcare professional immediately.

How should I store this medication?

Store in a cool and dry place, away from direct sunlight. Keep this medication away from children.

Record the date opened on the eye drop bottle. Discard this medication one month after opening.

How should I throw away this medication safely?

Pack this medication into a black trash bag and seal it tightly before throwing into the rubbish chute or bin.

Disclaimers

If you take more than the recommended dose, please seek medical advice immediately. The information provided on this page does not replace information from your healthcare professional. Please consult your healthcare professional for more information.

This article is jointly developed by members of the National Medication Information workgroup. The workgroup consists of cluster partners (National Healthcare Group, National University Health System and SingHealth), community pharmacies (Guardian, Unity and Watsons) and Pharmaceutical Society of Singapore. The content does not reflect drug availability and supply information in pharmacies and healthcare institutions. You are advised to check with the respective institutions for such information.

Last updated on Jun 2021

