

## PHARMACY

Drug Information

### FOLFIRI-Cetuximab Regimen [Fluorouracil, Leucovorin, Irinotecan, Cetuximab]



#### What is FOLFIRI-Cetuximab Regimen?

- FOLFIRI with Cetuximab is usually given for the treatment of colorectal cancer. It consists of four different drugs: Leucovorin (Folinic acid), Fluorouracil, Irinotecan and Cetuximab.
- Leucovorin is not a chemotherapy drug. It is often given with Fluorouracil, as it has been shown to make it works better.

#### How is FOLFIRI-Cetuximab Regimen Administered?

- FOLFIRI-Cetuximab is given as separate injections into your vein (intravenous) every 14 days.
- On the first day of treatment, you will receive Cetuximab, Irinotecan and Folinic acid as a drip (infusion). After this, you will have Fluorouracil given as an intravenous injection followed by an infusion over 46 hours via a small portable pump. You can go home once the infusion starts and return two days later for the nurse to disconnect your pump when your infusion is completed.

#### Can FOLFIRI-Cetuximab Regimen be Administered With Other Medications?

- Please inform your doctor or pharmacist if you are taking other medications, including Traditional Chinese Medicines, Jamu medicine or supplements.

#### What are the Important Side Effects of FOLFIRI-Cetuximab Regimen?

- Infusion related reactions/Allergic reactions  
These are usually mild or moderate, but rarely, they can be more severe. You may be given drugs before the infusion to reduce the chance of a reaction happening.

If you do have a reaction, this can usually be resolved by slowing or stopping the drip until the reaction is over. Any future infusions will be given at a slower rate.

*Inform the nurses immediately during infusion of FOLFIRI-Cetuximab if you experience the following:*

- Flu-like symptoms, such as headache, feeling flushed, having a fever, chills or dizziness
- Red, warm and itchy bumps on the skin
- Swelling in the lips, tongue or throat

- Breathlessness, wheezing, a cough or sudden difficulty of breathing
- Chest tightness or chest pain

Rarely, an infusion-related reaction can happen a few hours after treatment. If you develop any of the symptoms above or feel unwell after you get home, please proceed to the A & E department immediately.

- Pain along the vein  
FOLFIRI may cause pain at the site where the injection is given or along the vein. If you feel pain, tell your nurse straight away so that the infusion site can be checked. They may give the drug more slowly or flush it through with more fluids to reduce pain.
- Medications prescribed by your doctor should be beneficial for your condition even though they may result in some inevitable side effects.

Side Effect	Management
Risk of infection can occur as FOLFIRI can reduce the number of white blood cells in your blood	<ul style="list-style-type: none"> <li>• To reduce risk of infection:               <ul style="list-style-type: none"> <li>- Wash your hands after toileting.</li> <li>- Avoid crowds and people who are sick.</li> <li>- Avoid raw food.</li> </ul> </li> <li>• Please proceed to the A&amp;E department immediately if you experience signs of infection such as fever with a temperature of 38°C and above, chills and shivering. Avoid taking Paracetamol (Panadol®) to relieve fever. Be aware that taking Paracetamol may mask fever and hence delay treatment of infection.</li> </ul>
Bruising or bleeding can occur as FOLFIRI can reduce the number of platelets in your blood	<ul style="list-style-type: none"> <li>• To reduce risk of bleeding problems:               <ul style="list-style-type: none"> <li>- Do not pick your nose.</li> <li>- Brush your teeth gently with a soft toothbrush.</li> <li>- Be aware that bleeding may take a longer time to stop.</li> </ul> </li> <li>• Please proceed to the A&amp;E department immediately if you experience any severe bleeding (eg, passing a large amount of fresh blood in urine or stools, or passing black sticky stools) or significant bruising.</li> </ul>

Side Effect	Management
Nausea or vomiting	<ul style="list-style-type: none"> <li>• Take small, frequent meals.</li> <li>• Avoid smells that precipitate nausea.</li> <li>• Take medicines for nausea or vomiting prescribed by your doctor.</li> <li>• Avoid tight-fitting clothes around the waist.</li> </ul>
Early diarrhoea (may occur 24 hours after irinotecan treatment) and accompanied with watery eyes, stomach cramps and sweating.	<ul style="list-style-type: none"> <li>• Inform nurses/ doctor immediately if it happens while chemotherapy is being given or proceed to A &amp; E department if it happens when you get home.</li> </ul>
Late diarrhoea (may occur one day to several days after irinotecan treatment). It starts with stools more loose or often than usual.	<ul style="list-style-type: none"> <li>• Late diarrhea must be treated immediately with loperamide tablets.               <ul style="list-style-type: none"> <li>-Take two tablets at the first sign of loose or more frequent stools than usual.</li> <li>-Then take one tablet every two hours until diarrhea has stopped for 12 hours.</li> <li>-At night, you may take two tablets every four hours during the time you usually sleep.</li> <li>-This dose is much higher than the package directions. It is very important that you take this higher dose to stop the diarrhea.</li> <li>-Always keep a supply of loperamide (eg, have 48 tablets on hand). You can buy loperamide at any pharmacy without a prescription.</li> </ul> </li> <li>• Proceed to A &amp; E department if diarrhea does not improve 24 hours after starting loperamide or lasts more than 36 hours.</li> <li>• Drink at least two litres (8 cups) of fluids every day when you have diarrhoea.</li> <li>• Avoid milk or dairy products and/or high fibre food whilst you are having diarrhoea.</li> </ul>

Side Effect	Management
Mouth ulcers can occur on the tongue, the sides of the mouth or in the throat and this can lead to an infection	<ul style="list-style-type: none"> <li>Brush your teeth gently after eating and at bedtime with a soft toothbrush.</li> <li>Rinse and brush your dentures after eating and soak them in denture solution overnight.</li> <li>Rinsing your mouth regularly with an alcohol-free mouth gargle or salt water may help reduce chances of infection.</li> </ul>
Hand-foot syndrome may occur during treatment. The palms of hands and soles of your feet may tingle, become red, darken, numb, painful, swollen or creases may be observed. Skin may also become dry or itchy. Nail beds may also darken.	<ul style="list-style-type: none"> <li>Inform your doctor if blisters, severe pain or ulcers affects your normal daily activities.</li> <li>Avoid tight-fitting shoes or rubbing pressure to hands and feet if this occurs.</li> <li>Applying moisturizer to hands and feet liberally often may help.</li> <li>Applying sunscreen with an SPF (sun protection factor) of at least 30 may help.</li> </ul>
Acne-like rashes may appear on the face, upper chest and back usually within three weeks after starting treatment and go away completely when the treatment stops. Skin on the face may also become flaky and scaly.	<ul style="list-style-type: none"> <li>Avoid direct sunlight and applying sunscreen with an SPF (sun protection factor) of at least 30 may help. Alternatively, wear a hat, long sleeves, and long pants outside on sunny days.</li> <li>Applying moisturizers to dry areas may help.</li> </ul>
Nail changes, inflammation of the skin surrounding a fingernail or toenail may occur	<ul style="list-style-type: none"> <li>Mild pain can be relieved by taking Paracetamol (Panadol ®). Inform your doctor if pain is not controlled by Paracetamol (Panadol ®). Avoid Paracetamol (Panadol ®) when you are having fever.</li> <li>If you get pain, redness or swelling around your nails, inform your doctor.</li> </ul>

Side Effect	Management
Reduced levels of magnesium in blood	<ul style="list-style-type: none"> <li>Regular blood test is required to monitor magnesium levels in your blood and supplementation may be given if required.</li> </ul>
Hair Loss	<ul style="list-style-type: none"> <li>Wear a wig or hat or use a gentle shampoo.</li> </ul>
Fatigue or Tiredness	<ul style="list-style-type: none"> <li>Do not drive or operate machinery if you are feeling tired. Inform your doctor if fatigue is persistent or worsens.</li> </ul>
Heart problems	<ul style="list-style-type: none"> <li>Please proceed to the A &amp; E department immediately if you experience signs of heart problems such as shortness of breath, difficulty breathing, chest pain, fast heart beat or sudden swelling of legs.</li> </ul>

**If you develop rashes, facial swelling or shortness of breath after taking the medication, you could be allergic to the medication. Please seek medical attention immediately.**

#### Are There any Special Precautions That I Need to Take?

- It is important to inform your doctor if you intend to conceive or breastfeed while taking this medication.
- You must not become pregnant during treatment and must use an effective method of contraception. You should take appropriate contraceptive measures during and up to 6 months after cessation of treatment. If pregnancy occurs during your treatment, you must inform your doctor immediately.
- FOLFIRI-Cetuximab regimen may have an anti-fertility effect, which could be irreversible. You should discuss with your doctor regarding fertility preservation prior to starting treatment.
- You should not breast-feed during treatment.

#### What Else can I do to Help My Condition?

- Exercise

Light exercise is recommended as it helps to improve mobility, balance and even emotional well-being. Do avoid strenuous exercise, and rest when you are feeling tired.

- Stay positive

Understand your condition and take an active role in managing it. Do not be afraid to talk about it with your doctor/ pharmacist/ family members. Repressed feelings can be harmful to your immediate well-being and long-term health.

- Adequate hydration

Maintain adequate hydration with 2 to 3 litres (8 to 12 cups) of fluids per day unless informed by your doctor to restrict fluids. This will decrease your chances of being constipated and also prevent dehydration.

- Medical and dental treatment

If you need to go into hospital for any reason other than cancer, always inform the doctors and nurses that you are having chemotherapy.

Talk to your cancer doctor if you think you need dental treatment. Always inform your dentist that you are having chemotherapy

**If you have any problem with your treatment, please contact your doctor, pharmacist or nurse clinician.**

**You can also find useful information with regards to your medication on this website: <https://www.macmillan.org.uk/information-and-support>**

