

PHARMACY

Drug Information

Pertuzumab (Perjeta®) Injection



What is Pertuzumab?

Pertuzumab is usually used in combination with Trastuzumab for the treatment of breast cancer.

How is Pertuzumab Administered?

- Pertuzumab is usually given by injection into a vein (an intravenous infusion) over 30 to 60 minutes every three weeks.

Can Pertuzumab be Administered / Taken With Other Medications?

- Please inform your doctor or pharmacist if you are taking other medications, including Traditional Chinese Medicines, Jamu medicines or supplements.

What are the Important Side Effects of Pertuzumab?

Infusion related reaction

- These are usually mild or moderate, but rarely, they can be more severe. You may be given drugs before the infusion to reduce the chance of a reaction happening.
- If you do have an infusion related reaction, this can usually be treated by slowing or stopping the drip until the reaction is over. Any future infusions will be given at a slower rate.
- *Inform the nurses immediately during infusion of Pertuzumab if you experience the following:*
 - Flu-like symptoms, such as headache, feeling flushed, having a fever, chills or dizziness
 - Red, warm and itchy bumps on the skin
 - Swelling of the lips, tongue or throat
 - Breathlessness, wheezing, a cough or sudden difficulty breathing
 - Chest tightness or chest pain

- Medications prescribed by your doctor should be beneficial for your condition even though they may result in some inevitable side effects.

Side Effect	Management
Nausea or vomiting	<ul style="list-style-type: none"> • Take small, frequent meals. • Avoid smells that precipitate nausea. • Take medicines for nausea or vomiting prescribed by your doctor. • Avoid tight-fitting clothes around the waist.
Diarrhoea	<ul style="list-style-type: none"> • Inform your doctor if this is severe and/or does not improve with medicine use. • Drink at least 2 litres (8 cups) of fluids every day when you are having diarrhoea. • Avoid milk, dairy products, high fibre food or spicy food while you are having diarrhoea.
Muscle aches and joint pain	<ul style="list-style-type: none"> • Mild pain can be relieved by taking Paracetamol (Panadol ®). Inform your doctor if pain is not controlled by Paracetamol (Panadol ®). Avoid Paracetamol (Panadol ®) when you are having fever. • Taking regular warm baths may also help.
Changes the way the heart works	<ul style="list-style-type: none"> • You may require tests to check how well your heart is working before, during and sometimes after treatment. • Please proceed to the A & E department immediately if you experience shortness of breath, difficulty breathing, tightness in chest, chest pain, fast or irregular heart beat or sudden swelling of legs.

If you develop rashes, facial swelling or shortness of breath after taking the medication, you could be allergic to the medication. Please seek medical attention immediately.

Are There any Special Precautions That I Need to Take?

- It is important to inform your doctor if you intend to conceive or breastfeed while taking this medication.
- You must not become pregnant during treatment with Pertuzumab and must use an effective method of contraception. You should take appropriate contraceptive measures during and up to 7 months after stopping treatment. If pregnancy occurs during your treatment, you must inform your doctor immediately.
- Pertuzumab may have an anti-fertility effect, which could be irreversible. You should discuss with your doctor regarding fertility preservation prior to starting treatment.
- You should not breast-feed during treatment.

What Else can I do to Help My Condition?

- Exercise
Light exercise is recommended as it helps to improve mobility, balance and even emotional well-being. Avoid strenuous exercise, and rest when you are feeling tired.
- Stay positive
Understand your condition and take an active role in managing it. Do not be afraid to talk about it with your doctor/ pharmacist/ family members. Repressed feelings can be harmful to your immediate well-being and long-term health.
- Adequate hydration
Maintain adequate hydration with 2 to 3 liters (8 to 12 cups) of fluids per day unless informed by your doctor to restrict fluids. This will decrease your chances of getting constipation and also prevent dehydration.
- Medical and dental treatment
If you need to go into hospital for any reason other than cancer, always inform the doctors and nurses that you are having chemotherapy. Talk to your cancer doctor if you think you need dental treatment. Always inform your dentist that you are having chemotherapy.

If you have any problem with your treatment, please contact your doctor, pharmacist or nurse clinician.

You can also find useful information with regards to your medication on this website:

<https://www.macmillan.org.uk/information-and-support>



© Tan Tock Seng Hospital, Singapore 2021. All rights reserved. All information correct as of August 2021. No part of this document may be reproduced, copied, reverse compiled, adapted, distributed, commercially exploited, displayed or stored in a database, retrieval system or transmitted in any form without prior permission of Tan Tock Seng Hospital. All information and material found in this document are for purposes of information only and are not meant to substitute any advice provided by your own physician or other medical professionals.

PECC-PHA-ED-2021-1286-v1