

PHARMACY

Drug Information

Trifluridine-Tipiracil Capsule (Lonsurf®)



What is Trifluridine-Tipiracil?

- Trifluridine-Tipiracil is used for the treatment of metastatic stomach and colon cancer.

How is Trifluridine-Tipiracil Administered / Taken?

- Trifluridine-Tipiracil is usually taken orally twice a day within one hour after eating your morning and evening meals on Days 1 to 5 and Days 8 to 12 of a 28-day cycle. The doses should be spaced 10 to 12 hours apart.
- The capsule should be swallowed whole. Do not chew or crush the capsules. Inform your doctor or pharmacist if you have difficulty swallowing.
- If possible, handle the medication yourself. Otherwise, family member or caregiver can help to handle the medication. Wash hands before and after handling the medication.
- If you miss or vomit a dose of Trifluridine-Tipiracil, skip the missed dose and take it at the next scheduled timing.

Can Trifluridine-Tipiracil be Administered / Taken with Other Medications?

- Please inform your doctor or pharmacist if you are taking other medications, including Traditional Chinese Medicines, Jamu medicines or supplements.

What are the Important Side Effects of Trifluridine-Tipiracil?

- Medications prescribed by your doctor should be beneficial for your condition even though they may result in some inevitable side effects.

Side Effect	Management
Risk of infection can occur as Trifluridine-Tipiracil can reduce the number of white blood cells in your blood	<ul style="list-style-type: none"> • To reduce risk of infection: <ul style="list-style-type: none"> - Wash your hands after toileting. - Avoid crowds and people who are sick. - Avoid raw food. • Please proceed to the A&E department immediately if you experience signs of infection such as fever with a temperature of

	38°C and above, chills and shivering. Avoid taking Paracetamol (Panadol®) to relieve fever. Be aware that taking Paracetamol may mask fever and hence delay treatment of infection.
Bruising or bleeding can occur as Trifluridine - Tipiracil can reduce the number of platelets in your blood	<ul style="list-style-type: none"> To reduce risk of bleeding problems: <ul style="list-style-type: none"> - Do not pick your nose. - Brush your teeth gently with a soft toothbrush. - Be aware that bleeding may take a longer time to stop. Please proceed to the A&E department immediately if you experience any severe or persistent bleeding (e.g. passing a large amount of fresh blood in urine or stools, or passing black sticky stools) or significant bruising.
Nausea or vomiting	<ul style="list-style-type: none"> Take small, frequent meals. Avoid smells that precipitate nausea. Take medicines for nausea or vomiting prescribed by your doctor. Avoid tight-fitting clothes around the waist.
Diarrhoea	<ul style="list-style-type: none"> Inform your doctor if this is severe and/or does not improve with medicine use. Drink at least two litres (8 cups) of fluids every day when you are having diarrhoea. Avoid milk, dairy products, high fibre food or spicy food whilst you are having diarrhoea.
Fatigue or tiredness	<ul style="list-style-type: none"> Do not drive or operate machinery if you are feeling tired. Inform your doctor if fatigue is persistent or worsens.

If you develop rashes, facial swelling or shortness of breath after taking the medication, you could be allergic to the medication. Please seek medical attention immediately.

Are There any Special Precautions That I Need to Take?

- You must not become pregnant during treatment with Trifluridine-Tipiracil and must use an effective method of contraception. You should take appropriate contraceptive measures during and up to six months after stopping treatment. If pregnancy occurs during your treatment, you must inform your doctor immediately.
- Trifluridine-Tipiracil may have an anti-fertility effect, which could be irreversible. You should discuss with your doctor regarding fertility preservation prior to starting treatment.
- You should not breast-feed during treatment.

What Else Can I Do to Help My Condition?

- Exercise
Light exercise is recommended as it helps to improve mobility, balance and even emotional well-being. Avoid strenuous exercise, and rest when you are feeling tired.
- Stay positive
Understand your condition and take an active role in managing it. Do not be afraid to talk about it with your doctor/ pharmacist/ family members. Repressed feelings can be harmful to your immediate well-being and long-term health.
- Adequate hydration
Maintain adequate hydration with two to three litres (8 to 12 cups) of fluids per day unless informed by your doctor to restrict fluids. This will decrease your chances of getting constipation and also prevent dehydration.
- Medical and dental treatment
If you need to go into hospital for any reason other than cancer, always inform the doctors and nurses that you are receiving anti-cancer treatment.
Talk to your cancer doctor if you think you need dental treatment. Always inform your dentist that you are receiving anti-cancer treatment.

How Do I Store Trifluridine-Tipiracil?

- Store your medication in a cool, dry place away from heat, moisture and direct sunlight.

If you have any problem with your treatment, please contact your doctor, pharmacist or nurse clinician.

Please keep all medications out of reach of children.

You can also find useful information with regards to your medication on this website:

<https://www.macmillan.org.uk/information-and-support>



© Tan Tock Seng Hospital, Singapore 2021. All rights reserved. All information correct as of December 2021. No part of this document may be reproduced, copied, reverse compiled, adapted, distributed, commercially exploited, displayed or stored in a database, retrieval system or transmitted in any form without prior permission of Tan Tock Seng Hospital. All information and material found in this document are for purposes of information only and are not meant to substitute any advice provided by your own physician or other medical professionals.