

Eplerenone

Medication Information Leaflet

What is this medication for?

Eplerenone belongs to a class of medications called Mineralocorticoid Receptor Antagonist (MRA). It works on your kidneys to remove extra water and salts in your body. This will decrease the pressure on the blood vessels and allow the heart to work less hard. It is also used to reduce the amount of fluid present in the lungs as a result of heart failure.

Heart failure is a condition where the heart is weak and is unable to pump enough blood to the rest of the body. This medication is used in heart failure to improve the survival rate, reduce the symptoms of heart failure such as breathlessness or leg swelling and reduce the chance of being admitted to the hospital because of heart failure.

This medication may also be used to treat high blood pressure or high aldosterone (a type of hormone) levels.

How should I take/use this medication?

- Do not stop taking your medication without checking with your healthcare professional.
- You may take this medication with or without food.
- If you are taking one dose a day, take it in the morning. If you are taking the medicine two times a day, take the second dose before 6pm so that you do not need to wake up often at night to urinate.

What should I do if I forget to take/use this medication?

If you miss a dose, take the missed dose as soon as you remember. If it is almost time for your next dose, take only the usual dose. Do not double your dose or use extra medication to make up for the missed dose.

What precautions should I take?

Inform your healthcare professional if:

- You are allergic to this medication or any of the other ingredients of this medication
- You are pregnant, planning to become pregnant, or breastfeeding
- You have any kidney problems

What are some common side-effects of this medication?

Some side effects may include:

- Passing urine more often
- Dizziness and lightheadedness
 - This may occur especially when you get up quickly from lying or sitting down. Getting up slowly or changing posture slowly may help. Take a rest by sitting or lying down if you feel dizzy.
- Increased potassium levels in your blood
 - Your doctor may monitor your potassium levels and kidney function regularly when you are on this medication.

Inform your healthcare professional if these side effects do not go away and become bothersome to you.

What are some rare but serious side-effects that I need to seek medical advice immediately?

The symptoms of a drug allergy include one or more of the following:

- Swollen face/eyes/lips/tongue
- Difficulty in breathing
- Itchy skin rashes over your whole body

If you experience any of these symptoms, you should stop your medication and see your healthcare professional immediately.

What food or medication should I avoid when I take this medication?

Some medications may interact with eplerenone and affect their effectiveness or increase the risk of side-effects. However, these medications may also be important for your treatment. Inform your healthcare professional of the other medications you are currently taking. Examples include:

- Medications for bacterial infections (e.g. clarithromycin, erythromycin)
- Medications for fungal infections (e.g. itraconazole, ketoconazole)

How should I store this medication?

Store in a cool and dry place, away from direct sunlight. Keep this medication away from children. Throw away all expired medications.

How should I throw away this medication safely?

Pack this medication into a black trash bag and seal it tightly before throwing into the rubbish chute or bin.

Disclaimers

If you take more than the recommended dose, please seek medical advice immediately. The information provided on this page does not replace information from your healthcare professional. Please consult your healthcare professional for more information.

This article is jointly developed by members of the National Medication Information workgroup. The workgroup consists of cluster partners (National Healthcare Group, National University Health System and SingHealth), community pharmacies (Guardian, Unity and Watsons) and Pharmaceutical Society of Singapore.

Last updated on July 2019